

★-REDUCTIONS OF IDEALS AND PRÜFER V-MULTIPLICATION DOMAINS

E. HOUSTON, S. KABBAJ ^(*), AND A. MIMOUNI ^(*)

ABSTRACT. Let R be a commutative ring and I an ideal of R . An ideal $J \subseteq I$ is a reduction of I if $JI^n = I^{n+1}$ for some positive integer n . The ring R has the (finite) basic ideal property if (finitely generated) ideals of R do not have proper reductions. Hays characterized (one-dimensional) Prüfer domains as domains with the finite basic ideal property (basic ideal property). We extend Hays' results to Prüfer v -multiplication domains by replacing "basic" with "w-basic," where w is a particular star operation. We also investigate relations among $\star$ -basic properties for certain star operations $\star$.

INTRODUCTION

Throughout, all rings considered are commutative with identity. Let R be a ring and I an ideal of R . An ideal $J \subseteq I$ is a *reduction* of I if $JI^n = I^{n+1}$ for some positive integer n [15]. An ideal that has no reduction other than itself is called a *basic* ideal [7]. The notion of reduction was introduced by Northcott and Rees, who stated: "First, it defines a relationship between two ideals which is preserved under homomorphisms and ring extensions; secondly, what we may term the reduction process gets rid of superfluous elements of an ideal without disturbing the algebraic multiplicities associated with it" [15]. For both early and recent developments on reduction theory, we refer the reader to [10, 7, 8, 11, 15, 16, 17, 18].

In [7, 8], Hays investigated reductions of ideals in commutative rings with a particular focus on Prüfer domains. He studied the notion of basic ideal and examined domains subject to the basic ideal property (i.e., every ideal is basic). This class is shown to be strictly contained in the class of Prüfer domains (domains in which every nonzero finitely generated ideal is invertible); and a new characterization for Prüfer domains is provided; namely, a domain is Prüfer if and only if it has the finite basic ideal property (i.e., every finitely generated ideal is basic) [7, Theorem 6.5]. The second main result of these two papers characterizes domains with the (full) basic ideal property as one-dimensional Prüfer domains ([7, Theorem 6.1] combined with [8, Theorem 10]). Our primary goal is to extend Hays' results to Prüfer v -multiplication domains (PvMDs).

Let R be a domain and I a nonzero fractional ideal of R . The v - and t -closures of I are defined, respectively, by $I_v := (I^{-1})^{-1}$ and $I_t := \bigcup J_v$, where J ranges over the set of finitely generated subideals of I . Recall that I is a t -ideal if $I_t = I$ and a t -finite (or v -finite) ideal if there exists a finitely generated fractional ideal J of R such that $I = J_t = J_v$; and R is called a Prüfer v -multiplication domain (PvMD) if the set of its t -finite t -ideals forms a group under ideal t -multiplication $((I, J) \mapsto (IJ)_t)$. A useful characterization is that R is a PvMD if and only if each localization at a maximal t -ideal is a valuation domain [6, Theorem 5]. The class of PvMDs strictly contains the classes of factorial and Prüfer

Date: February 24, 2016.

2010 *Mathematics Subject Classification.* 13A15, 13A18, 13F05, 13G05, 13C20.

^(*) Supported by King Fahd University of Petroleum & Minerals under DSR Research Grant #: RG1328.

domains. The t -operation is nowadays a cornerstone of multiplicative ideal theory and has been investigated thoroughly by many commutative algebraists since the 1980's.

For the convenience of the reader, the following figure displays a diagram of implications summarizing the relations among many well-studied classes of domains, putting PvMDs in perspective. In the diagram, classes on top become the classes directly underneath by means of replacing the definitions with a corresponding t -version. For example, a GCD-domain is a domain in which I_t is principal for each nonzero finitely generated ideal I , and a PvMD is a domain in which each nonzero finitely generated ideal is t -invertible.




FIGURE 1. PvMDs in perspective

The t - and v -operations are examples of star operations (defined below). We also require the w -operation: for a nonzero ideal fractional I of a domain R , $I_w = \bigcup (I : J)$, where the union is taken over all finitely generated ideals J of R that satisfy $J_v = R$; equivalently, $I_w = \bigcap IR_M$, where the intersection is taken over the set of maximal t -ideals of R . It follows that for each I and maximal t -ideal M , we have $I_w R_M = IR_M$. (This can be done in greater generality—see [1].) In the diagram above, one can replace “ t ” by “ w ” to go from top to bottom.

In Section 1 we discuss the notion of $\star$ -basic ideals and prove that a domain with the finite $\star$ -basic ideal property ($\star$ -basic ideal property) must be integrally closed (completely integrally closed). We also observe that a domain has the v -basic ideal property if and only if it is completely integrally closed. Section 2 is devoted to generalizing Hays’ results; we show that a domain has the finite w -basic ideal property (w -basic ideal property) if and only if it is a PvMD (of t -dimension one). In Section 3, we present a diagram of implications among domains having various $\star$ -basic properties and give examples showing that most of the implications are not reversible. For example, a domain with the w -basic ideal property must also have the t -basic ideal property and a v -domain must have the finite v -basic ideal property, but neither implication is reversible.

Notation is standard, as in [5]. In particular, for a domain D with quotient field K and submodules A, B of K , we use $(A : B)$ to denote the D -module $\{x \in K \mid xB \subseteq A\}$.

1. ★-BASIC IDEALS

Let R be a domain with quotient field K , and let $\mathcal{F}(R)$ denote the set of nonzero fractional ideals of R . A map $\star : \mathcal{F}(R) \rightarrow \mathcal{F}(R)$, $I \mapsto I^\star$, is said to be a *star operation on R* if the following conditions hold for every nonzero $a \in K$ and $I, J \in \mathcal{F}(R)$: (1) $(aI)^\star = aI^\star$ and $R^\star = R$; (2) $I \subseteq I^\star$ and $I \subseteq J$ implies $I^\star \subseteq J^\star$; and (3) $I^{\star\star} = I^\star$. It is common to denote the trivial star operation ($I \mapsto I$) by “ d .”

Definition 1.1. Let R be an integral domain and $\star$ a star operation on R . Let I be a nonzero ideal of R .

- (1) An ideal $J \subseteq I$ is a $\star$ -reduction of I if $(JI^n)^\star = (I^{n+1})^\star$ for some integer $n \geq 0$. The ideal J is a *trivial $\star$ -reduction* of I if $J^\star = I^\star$.
- (2) I is $\star$ -basic if it has no $\star$ -reduction other than the trivial $\star$ -reduction(s).
- (3) R has the $\star$ -basic ideal property if every nonzero ideal (or, equivalently, every $\star$ -ideal) of R is $\star$ -basic.
- (4) R has the finite $\star$ -basic ideal property if every nonzero finitely generated ideal (or, equivalently, every $\star$ -finite ideal) of R is $\star$ -basic.

It is clear that $\star$ -reductions can be extended to fractional ideals; in particular, if R has the $\star$ -basic ideal property, then every nonzero fractional ideal of R is $\star$ -basic.

It is easy to see that if $\star_1 \leq \star_2$ are star operations on a domain R (meaning that $I^{\star_1} \subseteq I^{\star_2}$ for each $I \in \mathcal{F}(R)$), then each $\star_1$ reduction of an ideal is also a $\star_2$ -reduction. The converse is false. In particular, a t -reduction may not be a (d) -reduction. For a very simple example, let $R = k[x, y]$ be a polynomial ring in two indeterminates over a field k , and let $M = (x, y)$. Then M is basic, i.e., M has no reductions other than itself [7, Theorem 2.3]. On the other hand, $M_t = R$ (see, e.g., [13, Exercise 1, p. 102]), from which it follows that any power of M is a (trivial) t -reduction of M . (We give a “better” example following Proposition 1.4 below.)

Lemma 1.2. In an integral domain R , $\star$ -invertible ideals and $\star$ -idempotent ideals are $\star$ -basic.

Proof. Let $J \subseteq I$ be a $\star$ -reduction of the ideal I of R , so that $(JI^n)^\star = (I^{n+1})^\star$ for some positive integer n . If I is $\star$ -invertible, then multiplication by $(I^{-1})^n$ and taking $\star$ -closures immediately yields $J^\star = I^\star$. Next, assume that $(I^2)^\star = I^\star$. Then $I^\star = (I^{n+1})^\star = (JI^n)^\star \subseteq J^\star \subseteq I^\star$ so that, again, $J^\star = I^\star$, as desired. $\square$

Lemma 1.3. (cf. [7, Lemma 6.4]) Let $\star$ be a star operation on a domain R . If R has the finite $\star$ -basic ideal property, then R is integrally closed.

Proof. Let $x, y \in R$ be such that x/y is integral over R . As in the proof of [7, Lemma 6.4], (y) is a reduction of (x, y) . We then have $x \in (x, y)^\star = (y)^\star = (y)$, whence $x/y \in R$. $\square$

Recall that a domain R is said to be *completely integrally closed* if every nonzero ideal of R is v -invertible.

Proposition 1.4. Let $\star$ be a star operation on an integral domain R .

- (1) If R has the $\star$ -basic ideal property, then R is completely integrally closed.
- (2) R has the v -basic ideal property if and only if R is completely integrally closed.

Proof. (1) Assume R has the $\star$ -basic ideal property. Let I be a nonzero ideal of R and set $J := II^{-1}$. It is well known that $J^{-1} = (J : J)$, and hence J^{-1} is a ring. Now, let $0 \neq a \in J$ and set $A := aJ^{-1}$ and $B := aR$. Clearly, A and B are v -ideals of R with $B \subseteq A$ and $BA = A^2$.

That is, B is a reduction (and, a fortiori, a $\star$ -reduction) of A . By the $\star$ -basic hypothesis, $aJ^{-1} = A^* = B^* = aR$, whence $R = J^{-1}$. Therefore, $(II^{-1})_v = J_v = R$, as desired.

(2) The “only if” assertion is a special case of (1), and the converse is handled by Lemma 1.2. $\square$

Next, we give an example of t -ideals I, J in a Noetherian domain R such that J is a t -reduction, but not a d -reduction, of I . Since the v - and t -operations coincide in any Noetherian domain, such an R cannot be (completely) integrally closed by Proposition 1.4.

Example 1.5. Again let k be a field and x, y indeterminants over k . Let $T = k[x, y] = k + M$, where $M = (x, y)T$. Now let $R = k + M^2$. Observe that R is Noetherian (see, e.g., [2]). As in the discussion preceding Lemma 1.2, as an ideal of T , M has no reductions other than itself. In particular, M^2 is not a reduction of M in T , and it follows easily that M^2 is not a reduction of (the fractional ideal) M in R . However, we claim that M^2 is a nontrivial t -reduction of M . To verify this, proceed as follows. First, we have $(T : M) = T$ (as before). It follows that $M \subseteq M^{-1} (= (R : M)) \subseteq T$. On the other hand, if $f \in T$ satisfies $fM \subseteq R$, then, writing $f = a + m$ with $a \in k$ and $m \in M$, we immediately obtain that $aM \subseteq R$, whence $a = 0$, i.e., $f \in M$. Thus $M^{-1} = M$, whence also $M_t = M_v = M$. However, $(R : T) = M^2$, whence $(M^2)^{-1} = ((R : M) : M) = (M : M) = T$ and then $(M^2)_t = (M^2)_v = (R : T) = M^2$. A similar argument yields $(M^n)_t = M^2$ for $n \geq 2$. Hence $M^2 = (M^3)_t = (M^2M)_t$, and therefore $J := M^2$ is a nontrivial t -reduction of $I := M$, as claimed. (To obtain an example involving integral ideals, replace M by xM and M^2 by xM^2 .) $\square$

We recall that a domain R is a v -domain if each nonzero finitely generated ideal of R is v -invertible. From Lemma 1.2, the following is immediate:

Proposition 1.6. *A v -domain has the finite v -basic ideal property.* $\square$

Now recall that to any star operation $\star$ on a domain R , we may define an associated star operation $\star_f$ by setting, for each $I \in \mathcal{F}(R)$, $I^{\star_f} = \bigcup J^\star$, the union being taken over all finitely generated subideals J of I ; the star operation $\star$ has *finite type* if $\star = \star_f$. Note that $v_f = t$. If $\star$ is a finite-type star operation on a domain R , then minimal primes of $\star$ -ideals are themselves $\star$ -ideals and each $\star$ -ideal is contained in a maximal $\star$ -ideal.

Lemma 1.7. *Let $\star$ be a star operation of finite type on an integral domain R . If I is a finitely generated ideal of R and J is a $\star$ -reduction of I , then there is a finitely generated ideal $K \subseteq J$ such that K is a $\star$ -reduction of I .*

Proof. Suppose that I is a finitely generated ideal of R and that $(JI^n)^\star = (I^{n+1})^\star$ for some ideal $J \subseteq I$ and some positive integer n . Suppose that I^{n+1} is generated by $b_1, \dots, b_r$ in R . Since $b_i \in (JI^n)^\star$, there is a finitely generated subideal K_i of J such that $b_i \in (K_i I^n)^\star$. For $K = \sum_{i=1}^r K_i$, we then have $I^{n+1} \subseteq (KI^n)^\star$, as desired. $\square$

Proposition 1.8. *If a domain R has the finite $\star$ -basic ideal property, then R also has the finite $\star_f$ -basic ideal property. In particular, if R has the finite v -basic ideal property, then R also has the finite t -basic ideal property.*

Proof. Let R be a domain with the $\star$ -basic ideal property. Let I be a finitely generated ideal of R , and let J be a $\star_f$ -reduction of I . By Lemma 1.7 we may assume that J is finitely generated. Since J is also a $\star$ -reduction of I , we have $J^{\star_f} = J^\star = I^\star = I^{\star_f}$. Hence R has the $\star_f$ -basic ideal property. $\square$

Corollary 1.9. *A v -domain has the finite t -basic ideal property.* $\square$

2. CHARACTERIZATIONS

We begin with an analogue of Hays' first result that a domain is a Prüfer domain if and only if it has the finite basic ideal property. We shall need a result of Kang [12, Theorem 3.5] that characterizes PvMDs as integrally closed domains in which the t - and w -operations coincide. We denote the set of maximal t -ideals of a domain R by $\text{Max}_t(R)$.

Theorem 2.1. (cf. [7, Theorem 6.5]) *A domain R is a PvMD if and only if it has the finite w -basic ideal property.*

Proof. If R is a PvMD, then, as mentioned above, the t - and w -operations coincide, and R has the finite w -basic ideal property by Corollary 1.9.

Now assume that R has the finite w -basic ideal property. Then R is integrally closed by Lemma 1.3. Let $M \in \text{Max}_t(R)$, and let $a, b \in M$. Since (a^2, b^2) is a reduction of $(a, b)^2$, we have $(a^2, b^2)_w = ((a, b)^2)_w$ and hence (as mentioned in the introduction) $(a^2, b^2)R_M = (a, b)^2R_M$. Thus R_M is a valuation domain [5, Theorem 24.3(4)]. Therefore, R is a PvMD. $\square$

Hays proved that, in a Prüfer domain, the definition of a reduction can be restricted; namely, $J \subseteq I$ is a reduction if and only if $JI = I^2$ [8, Proposition 1]. The next lemma establishes a similar property for t -reductions and also shows that this notion is local in the class of PvMDs. It is useful to note if J is a t -reduction of an ideal I , then a prime t -ideal of R contains I if and only if it contains J . We shall also need the fact (which follows easily from [19, Lemma 4] and is stated explicitly in [12, Lemma 3.4]), that if I is a nonzero ideal of a domain R and S is a multiplicatively closed subset of R , then $(I_t R_S)_{t_{R_S}} = (IR_S)_{t_{R_S}}$.

Lemma 2.2. *Let R be a PvMD and $J \subseteq I$ nonzero ideals of R . Then, the following assertions are equivalent:*

- (1) J is a t -reduction of I ;
- (2) $JR_M IR_M = (IR_M)^2$ for each $M \in \text{Max}_t(R)$;
- (3) $(JI)_t = (I^2)_t$.

Proof. (1) $\Rightarrow$ (2) Assume that J is a t -reduction of I , so that $(JI^n)_t = (I^{n+1})_t$ for some positive integer n , and let $M \in \text{Max}_t(R)$. Since R_M is a valuation domain, the t -operation is trivial on R_M ($t_{R_M} = d_{R_M}$). Using this and the remarks above, we have

$$I^{n+1}R_M = ((I^{n+1})_t R_M)_{t_{R_M}} = ((JI^n)_t R_M)_{t_{R_M}} = JI^n R_M.$$

Hence JR_M is a t -reduction of IR_M in R_M , and so $JR_M IR_M = (IR_M)^2$ by [8, Proposition 1].

(2) $\Rightarrow$ (3) By [12, Theorem 3.5], we have

$$(JI)_t = \bigcap_{M \in \text{Max}_t(R)} JIR_M = \bigcap_{M \in \text{Max}_t(R)} (I^2 R_M) = (I^2)_t.$$

(3) $\Rightarrow$ (1) is trivial. $\square$

Lemma 2.3. (cf. [8, Lemma 9]) *Let x be a nonzero element of a PvMD R , let P be a minimal prime of xR , and let $I = xR_P \cap R$. Then*

- (1) I is a w -ideal of R ,
- (2) $xR + I^2$ is a w -reduction of I , and
- (3) if I is w -basic, then $P \in \text{Max}_t(R)$.

Proof. (1)-(2) Let M be a maximal t -ideal of R containing P . Then $I_w \subseteq IR_M \cap R \subseteq IR_P \cap R = I$, proving (1). We next claim that $IR_M = IR_P \cap R_M$. To see this, suppose that $y \in IR_P \cap R_M$. Then we may write $y = a/s = b/t$ with $a \in I$, $b \in R$, $s \in R \setminus P$ and $t \in R \setminus M$.

We then have $b = at/s \in IR_P \cap R = I$, and hence $y = b/t \in IR_M$, as desired. Now, for $s \in R \setminus P$ and $a \in I$ (using the fact that R_M is a valuation domain), it is clear that $a/s \in IR_P \cap R_M = IR_M$. If we also have $b \in I$, then, writing $b = x/s'$ with $s' \in R \setminus P$, we obtain $ab = (a/s')x \in xIR_M$. Thus $I^2R_M = xIR_M$, and it follows that $(xR + I^2)IR_M = xIR_M = I^2R_M$. (In particular, $I^2R_M \subseteq xR_M$; we use this below.) Since I is P -primary, we also have $(xR + I^2)IR_N = I^2R_N$ for $N \in \text{Max}_t(R)$ with $N \not\subseteq P$. Therefore, $((xR + I^2)I)_w = (I^2)_w$, and so $xR + I^2$ is a w -reduction of I .

(3) Assume that I is w -basic; then $(xR + I^2)_w = I_w$ by (2). Suppose that $M \in \text{Max}_t(R)$ properly contains P , and choose $y \in M \setminus P$. Then P is minimal over yx , and $I = yxR_P \cap R$. Hence, as above, we have (using the parenthetical “in particular” comment above) $xR_M \in IR_M = (yxR + I^2)R_M \subseteq yxR_M$, a contradiction. Therefore, $P \in \text{Max}_t(R)$. $\square$

Theorem 2.4. *A domain R has the w -basic ideal property if and only if R is a PvMD of t -dimension 1.*

Proof. Let R be a PvMD with $t\text{-dim}(R) = 1$, and let $J \subseteq I$ be a nonzero ideals of R with $(JI)_w = (I^2)_w$. Let M be a maximal t -ideal of R . Then $JIR_M = I^2R_M$. We wish to show that $JR_M = IR_M$, and for this we may as well assume that $I \subseteq M$ and IR_M is not invertible. Since R_M is a valuation domain, we then have $IR_M = IMR_M$, and since R_M is also one-dimensional, [3, Proposition 2.1] yields $IR_M(R_M : IR_M) = MR_M$. Hence multiplying both sides of the equation $JIR_M = I^2R_M$ by $(R_M : IR_M)$ yields $JR_M \supseteq JMR_M = IMR_M = IR_M$. We then obtain $J_w = I_w$. Therefore, by Lemma 2.2, R has the w -basic ideal property.

Conversely, suppose that R has the w -basic ideal property. Then R is a PvMD by Theorem 2.1. Let M be a maximal t -ideal of R , let Q be a nonzero prime of R contained in M , let x be a nonzero element of Q , and shrink Q to a prime P minimal over x . Then, since $I := xR_P \cap R$ is w -basic by hypothesis, Lemma 2.3 yields $P = Q = M$. Therefore, $\text{ht} M = 1$, as desired. $\square$

3. EXAMPLES.

Consider the following diagram of implications involving various $\star$ -basic properties.

Of these implications, (1)-(3) and (9) are well known. Implications (4)-(8) follow from Proposition 1.6, Proposition 1.8, Lemma 1.3, Theorem 2.4 (and the fact that $w = t$ in a PvMD), and Proposition 1.4, respectively.

Irreversibility of arrows (1)-(3) and (9) is again well known. We do not know whether (5) is reversible. The remainder of the paper is devoted to examples for (irreversibility of) the other implications.

Example 3.1. Arrow (4) is irreversible.

Proof. Let k be a field and X, Y, Z indeterminates over k . Let $T := k((X)) + M$ and $R := k[[X]] + M$, where $M := (Y, Z)k((X))[[Y, Z]]$. Let A be an ideal of R . Then A is comparable to M . Suppose $A \subseteq M$ and A is not invertible. If $AA^{-1} \supsetneq M$, then AA^{-1} is principal, and hence A is invertible, contrary to assumption. Hence $AA^{-1} \subseteq M$. We claim that $(AA^{-1})_v = M$. To verify this, first recall that M is divisorial in R . Then, since AA^{-1} is a trace ideal, that is, $(AA^{-1})^{-1} = (AA^{-1} : AA^{-1})$, we have $(AA^{-1})^{-1} \subseteq (AA^{-1}T : AA^{-1}T) = T = M^{-1}$ (the first equality holding since T is Noetherian and integrally closed). This forces $(AA^{-1})^{-1} = M^{-1}$, whence $(AA^{-1})_v = M_v = M$, as claimed. Now let I be a finitely generated ideal of R and J a v -reduction of I , so that $(JI^n)_v = (I^{n+1})_v$ for some positive integer n . We shall show that $J^{-1} = I^{-1}$ (and hence that $J_v = I_v$), and for this we may assume that I is not invertible. Suppose, by way of contradiction, that $IT(T : IT) = T$, i.e., that IT is invertible




FIGURE 2. ★-basic properties in perspective

in T . Then, since T is local, IT is principal and, in fact, $IT = aT$ for some $a \in I$. We then have $R \subseteq a^{-1}I \subseteq T$. Then $k[[X]] \cong R/M \subseteq a^{-1}I/M \subseteq T/M \cong k((X))$, from which it follows that $a^{-1}I/M$ must be a cyclic $k[[X]]$ -module. However, this is easily seen to imply that $a^{-1}I$, hence I , is principal, the desired contradiction. We therefore have $(T : IT)I \subseteq M$, whence

$$(IM)^{-1} = (R : IM) = ((R : M) : I) = (T : I) = (M : I) \subseteq I^{-1}.$$

This immediately yields $I^{-1} = (IM)^{-1}$.

Now set $Q = I^n(I^n)^{-1}$. From above (setting $A = I^n$), we have $Q_v = M$. Therefore,

$$I^{-1} \subseteq J^{-1} \subseteq (JM)^{-1} = (JQ)^{-1} = (IQ)^{-1} = (IM)^{-1} = I^{-1},$$

which yields $J^{-1} = I^{-1}$, as desired. Hence R has the finite v -basic property. Finally, again from above, we have $((y, z)(y, z)^{-1})_v = M$, so that R is not a v -domain. □

Example 3.2. Arrow (6) is irreversible.

Proof. Let k be a field and X, Y indeterminates over k . Let $V = k(X)[[Y]]$ and $R = k + M$, where $M = Yk(X)[[Y]]$. Clearly, R is an integrally closed domain. Of course, M is divisorial in R . Also, $(M^2)^{-1} = ((R : M) : M) = (V : M) = Y^{-1}V$, and so $(M^2)_v = (R : Y^{-1}V) = Y(R : V) = YM = M^2$, i.e., M^2 is also divisorial. We claim that R does not have the finite t -basic ideal property. Indeed, let $W := k + Xk$ and consider the finitely generated ideal I of R given by $I = Y(W + M)$. We have $(k : W) = (0)$; otherwise, we have $0 \neq f \in (k : W)$, and both f and $fX \in k$, whence $X \in k$, a contradiction. Therefore, $I^{-1} = Y^{-1}M$ and thus $I_t = I_v = YM^{-1} = M$. Now, let $J = YR$. Then $J_t = YR \subsetneq M = I_t$. However,

$$(JI)_t = (YI)_t = YI_t = YM = M^2 = ((I_t)^2)_t = (I^2)_t,$$

and so R does not have the finite t -basic ideal property. □

Example 3.3. Arrow (7) is irreversible.

Proof. In [9] Heinzer and Ohm give an example of an essential domain that is not a PvMD. In that example, k is a field, y, z , and $\{x_i\}_{i=1}^\infty$ are indeterminates over k , and $D = R \cap (\bigcap_{i=1}^\infty V_i)$, where $R = k(\{x_i\})[y, z]_{(y,z)k(\{x_i\})[y,z]}$ and V_i is the rank-one discrete valuation ring on $k(\{x_j\}_{j=1}^\infty, y, z)$ with x_i, y, z all having value 1 and x_j having value 0 for $j \neq i$ (using the “infimum” valuation). As further described in [14, Example 2.1], we have $\text{Max}(D) = \{M\} \cup \{P_i\}$, where M is the contraction of $(y, z)R$ to D and the P_i are the centers of the maximal ideals of the V_i ; moreover, $D_M = R$ and $V_i = D_{P_i}$.

It was pointed out in [4, Example 1.7] that each finitely generated ideal of D is contained in almost all of the V_i . In fact, one can say more. Let a be an element of D . We may represent a as a quotient f/g with $f, g \in T := k[\{x_i\}, y, z]_{(y,z)k[\{x_i\}, y, z]}$ and $g \notin (y, z)T$ (and hence $g \notin M$). Since f and g involve only finitely many x_j and $g \notin M$, the sequence $\{v_i(a)\}$ must be eventually constant, where v_i is the valuation corresponding to V_i . We denote this constant value by $w(a)$. A similar statement holds for finitely generated ideals of D .

Let K be a nonzero ideal of D . Then

$$K_t D_{P_i} \supseteq K D_{P_i} = (K D_{P_i})_{t D_{P_i}} = (K_t D_{P_i})_{t D_{P_i}} \supseteq K_t D_{P_i},$$

whence $K_t D_{P_i} = K D_{P_i}$.

Now suppose that we have nonzero ideals $J \subseteq I$ of D with $(J I^n)_t = (I^{n+1})_t$. Let $a \in I$, and choose $a_0 \in I$ so that $w(a_0)$ is minimal. Then $a a_0^n \in I^{n+1} \subseteq (J I^n)_t$, and so $a a_0^n \in (B A^n)_v$ for finitely generated ideals $B \subseteq J$ and $A \subseteq I$. With the observation in the preceding paragraph, we then have $a a_0^n \in B A^n D_{P_i}$ for each i . However, since $w(a_0) \leq w(A)$, it must be the case that $w(a) \geq w(B)$; i.e., for some integer k , $a \in B D_{P_i}$ for all $i > k$. Since the equality $(J I^n)_t = (I^{n+1})_t$ yields $J D_{P_i} = I D_{P_i}$ for each i , we may choose elements $b_j \in J$ for which $v_j(a) = v_j(b_j)$, $j = 1, \dots, k$. With $B' = (B, b_1, \dots, b_k)$, we then have $a \in B' D_{P_i}$ for each i . This yields $a(B')^{-1} \subseteq \bigcap D_{P_i}$.

Next, we consider extensions to D_M . From $(J I^n)_t = (I^{n+1})_t$, we obtain $(J I^n D_M)_{t D_M} = (I^{n+1} D_M)_{t D_M}$. Since D_M is a regular local ring, each nonzero ideal of D_M is t -invertible, and we may cancel to obtain $(I D_M)_{t D_M} = (J D_M)_{t D_M}$. There is a finitely generated subideal B_1 of J with $B_1 D_M = J D_M$. We then have

$$I B_1^{-1} \subseteq I D_M B_1^{-1} D_M = I D_M (B_1 D_M)^{-1} \subseteq (J D_M (J D_M)^{-1})_{t D_M} \subseteq D_M.$$

Now let $B_2 = B' + B_1$. Then $a(B_2)^{-1} \subseteq D_M \cap \bigcap D_{P_i} = D$, whence $a \in (B_2)_v \subseteq J_t$. It follows that D has the t -basic property. However, since D is not a PvMD, D cannot have the (finite) w -basic property. $\square$

Example 3.4. Arrow (8) is irreversible.

Proof. Let D denote the ring of entire functions. It is well known that D is a completely integrally closed Prüfer domain of infinite Krull dimension. Since D is a Prüfer domain, each nonzero ideal is a t -ideal. The fact that $\dim D = \infty$ then yields that D does not have the $(t-)$ basic property by [8, Theorem 10]. $\square$

REFERENCES

- [1] D.D. Anderson and S. J. Cook, *Two star-operations and their induced lattices*, Comm. Algebra **28** (2000), 2461-2475.
- [2] J. Boynton and S. Sather-Wagstaff, *Regular pullbacks*, In: Progress in Commutative Algebra 2 145-169, Walter de Gruyter, Berlin, 2012.

- [3] M. Fontana, J. Huckaba, and I. Papick, *Domains satisfying the trace property*, J. Algebra **107** (1987) 169–182.
- [4] S. Gabelli, E. Houston, and T. Lucas, *The t -# property for integral domains*, J. Pure Appl. Algebra **194** (2004), 281–298.
- [5] R. Gilmer, *Multiplicative ideal theory*, Dekker, New York, 1972.
- [6] M. Griffin, *Some results on v -multiplication rings*, Canad. J. Math. **19** (1967), 710–722.
- [7] J. Hays, *Reductions of ideals in commutative rings*, Trans. Amer. Math. Soc. **177** (1973) 51–63.
- [8] J. Hays, *Reductions of ideals in Prüfer domains*, Proc. Amer. Math. Soc. **52** (1975) 81–84.
- [9] W. Heinzer and J. Ohm, *An essential ring which is not a v -multiplication ring*, Can. J. Math. **25** (1973), 856–861.
- [10] W. Heinzer, L. Ratliff, and D. Rush, *Reductions of ideals in local rings with finite residue fields*, Proc. Amer. Math. Soc. **138** (5) (2010) 1569–1574.
- [11] C. Huneke and I. Swanson, *Integral closure of ideals, rings, and modules*, London Mathematical Society Lecture Note Series, 336. Cambridge University Press, Cambridge, Online version, 2012.
- [12] B. G. Kang, *Prüfer v -multiplication domains and the ring $R[X]_{N_v}$* , J. Algebra **123** (1989), 151–170.
- [13] I. Kaplansky, *Commutative rings*, revised edition, Univ. Chicago Press, Chicago, 1974.
- [14] J. L. Mott and M. Zafrullah, *On Prüfer v -multiplication domains*, Manuscripta Math. **35** (1981) 1–26.
- [15] D. G. Northcott and D. Rees, *Reductions of ideals in local rings*, Proc. Cambridge Philos. Soc. **50** (1954) 145–158.
- [16] D. Rees and J. Sally, *General elements and joint reductions*, Michigan Math. J. **35** (1988) 241–254.
- [17] Y. Song and S. Kim, *Reductions of ideals in commutative Noetherian semi-local rings*, Commun. Korean Math. Soc. **11** (3) (1996) 539–546.
- [18] H.-J. Wang, *Core of ideals of Noetherian local rings*, Proc. Amer. Math. Soc. **136** (3) (2008) 801–807.
- [19] M. Zafrullah, *On finite-conductor domains*, Manuscripta Math. **24** (1978), 191–204.

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF NORTH CAROLINA AT CHARLOTTE,
CHARLOTTE, NC 28223, USA

E-mail address: eghousto@uncc.edu

DEPARTMENT OF MATHEMATICS AND STATISTICS, KING FAHD UNIVERSITY OF PETROLEUM & MIN-
ERALS, DHAHRAN 31261, KSA

E-mail address: kabbaj@kfupm.edu.sa

DEPARTMENT OF MATHEMATICS AND STATISTICS, KING FAHD UNIVERSITY OF PETROLEUM & MIN-
ERALS, DHAHRAN 31261, KSA

E-mail address: amimouni@kfupm.edu.sa